

**Evaluation Summary of
Carob bean extract
for Use as a Cigarette Ingredient**

October 2005

INTRODUCTION

Carob bean extract (CAS # 84961-45-5) is currently used worldwide at levels below **5,000 ppm** in selected cigarette brands manufactured and/or distributed by Philip Morris International. Carob bean extract may be applied directly to the tobacco during cigarette manufacturing as a tobacco flavoring material, and as such, may be subject to pyrolysis-type reactions. Carob bean extract may also be applied to the filter as a flavoring material where it would not be subjected to pyrolysis temperatures. This document summarises our internal studies and current published toxicology information on carob bean extract abstracted from online toxicity databases.

TOXICITY DATA ON UN-BURNED MATERIAL

Overview

Carob bean extract is Generally Recognized As Safe (GRAS) by the U.S. Food and Drug Administration for direct food additive in accordance with 21 CFR § 182.20.¹ The Flavor and Extract Manufacturers Association (FEMA No. 2243)² and the Council of Europe (CoE No. 120)³ also approve carob bean extract for food use.

Carob bean extract is a product delivered in a tincture of alcohol (generally 20-25% carob in 55-77% alcohol) or other fluids.⁴ The extract is derived from milled seeds of the fruit pod of *Ceratonia siliqua Leguminosae*, a large evergreen tree native to the Middle East region and to the southern part of the Mediterranean. It is extensively cultivated for the seeds and fruit in Cyprus, Greece, Syria, Spain and Italy.⁵ The carob tree (also referred to as the locust tree) can reach heights of 33 feet and is well adapted to arid farming conditions.⁶

Health hazard data

A thorough search of the literature on carob bean failed to reveal significant information regarding the potential toxicity of carob bean extract. Since carob bean gum is used extensively in food manufacturing, most of the relevant published literature data involved studies conducted with carob bean gum. This summary therefore includes data derived from studies with carob bean gum. The relevance of the results of testing with carob bean gum to those that might be produced by carob bean extract is unknown.⁴

Carob bean gum is chemically defined as a 1,4-galactomannan polymer and biologically characterized as an indigestible polysaccharide hydrocolloid.⁷ Carob bean gum moves through the small intestine and arrives in the lower bowel largely undigested.⁸⁻¹⁰ Although decreases in serum glucose and cholesterol have been associated with ingestion of large amounts of carob bean gum by experimental animals,¹¹⁻¹⁴ the research information speaks to a secondary mechanism on these findings – through a reduction in gastric emptying and perhaps peristalsis, thereby delaying or retarding their absorption.¹³

The toxicology studies with carob bean gum further support the inert nature of this molecule. Acute toxicity studies in several species did not show any evidence of oral toxicity at doses up to 10 g/kg.¹⁵⁻¹⁷ Subchronic toxicity studies of dose levels in rodents up to 10% were unremarkable.¹⁶⁻¹⁸ Similarly, dogs receiving up to 10% (7,500 mg/kg/day) of carob bean gum admixed in the diet for 30 weeks showed no evidence of a compound-related toxicity.¹⁹ Lifetime bioassay studies in rats and mice with up to 5% (2,500 mg/kg/day for rats and 7,500 mg/kg/day for mice) dietary exposure to carob bean gum did not evoke any meaningful evidence that galactomannan possessed a carcinogenic (or a long-term toxicological) potential.^{16,20} Mutagenicity studies were negative^{17,21-23} and no chromosomal damage has

been demonstrated.^{17,23} Teratology studies performed in four species (rats,²⁴ mice,²⁴ hamsters²⁵ and rabbits²⁵) gave no suggestion of a teratogenic effect at levels of 910–1,300 mg/kg, although maternal toxicity was evident in rabbits and mice at the highest doses.^{24,25}

Carob bean gum has similarly low toxicity in humans. Several clinical studies have failed to show that oral ingestion of carob bean gum has any adverse sequelae.^{12,26,27} There is clinical data to support claims of a decrease in serum cholesterol and a blunting of the glucose-tolerance test following oral ingestion of carob bean gum, but whether this is directly related to a carob bean gum effect on gastrointestinal peristalsis or a consequence of the physical nature of the hydrocolloid structure is not known.^{12,28-31} The impact of these serum biochemical findings has not been readily embraced by the medical community because carob bean gum is not normally a recommended dietary approach to correcting even marginal increases in plasma cholesterol levels or fasting glucose swings.⁴

The only finding of consequence occurred with carob bean flour, a material derived from the unrefined whole ground seed. There are reports of an allergic response to extensive inhalation exposure to this flour.³²⁻³⁵ It is not known if this response is a rare occurrence. However, it does follow a pattern that has been noted with inhalation exposure to other proteinaceous materials, including soy protein.⁴

TOXICITY DATA ON BURNT MATERIAL

Combustion studies

As suggested by the purge and trap studies conducted by PM USA,³⁶ carob bean extract would not be expected to extensively distill at 100°C. Pyrolysis analysis at temperatures up to 900°C suggested that carob bean extract would pyrolyze extensively.³⁷

Philip Morris ingredient testing program

Carob bean extract was a part of the PM USA ingredient testing program that was designed to evaluate the potential effects of ingredients added to typical commercial blended test cigarettes on selected biological and chemical endpoints. Carob bean extract was added to test cigarette tobacco at target concentrations of 1,500 ppm, 4,500 ppm, or 7,000 ppm, and did not increase the mutagenic response of Salmonella bacteria to smoke condensate preparations.³⁸ Similarly, at the same target concentrations, the cytotoxic response of mouse embryo BALB/c 373 cells treated with mainstream smoke condensate preparations (particulate and gas vapor phase) was not altered by carob bean extract addition.³⁹ There were also no effects on any smoke constituents which correlated with the level of carob bean extract added to the cigarette tobacco on an equal TPM or tar basis.⁴⁰ The biological effects of inhaling smoke from cigarettes with carob bean extract was assessed in Sprague-Dawley rats exposed nose-only to smoke for 6 hrs/day, 7 days/week for 13 weeks. The results of the smoke inhalation studies indicated that the addition of carob bean extract to cigarette tobacco at target levels up to 7,000 ppm did not discernibly alter the biological effects normally associated with smoke exposure in rodents.⁴¹

CONCLUSION

Cigarette smoking causes lung cancer, heart disease, emphysema and other serious diseases in smokers. Smokers are far more likely to develop serious diseases, like lung cancer, than non-smokers. There is no "safe" cigarette. Government health warnings about smoking apply to all cigarettes, regardless of the ingredients added, including those containing only tobacco and paper.

The results of this evaluation of carob bean extract involving a review of published information and internal studies show there is no meaningful difference in the composition or toxicity of smoke from cigarettes with carob bean extract compared to the smoke from cigarettes without carob bean extract.³⁸⁻⁴¹ It is our scientific judgement, based on the best available data, that carob bean extract used in our cigarettes does not increase the overall toxicity of cigarette smoke.

References

1. Food and Drug Administration (1974) Part 121 - Food Additives. Subpart B - Exemption of certain food additives from the requirement of tolerances. Locust (carob) bean gum. Report No. Federal Register 39. p.34181-34184.
2. Hall, R.L. and Oser, B.L. (1965) Recent progress in the consideration of flavoring ingredients under the Food Additives Amendment: III.GRAS Substances. *Food Tech.* 253(2):151-197.
3. CoE (1981) *Flavouring Substances and Natural Sources of Flavourings*. Council of Europe Publishing, F-67006 Strausbourg Cedex, France. p.59-113.
4. Burdock, G.A. (2000) Safety Assessment of Carob Bean Gum as a Food Ingredient. Burdock Group, Vero Beach, FL. Unpublished Review.
5. LSRO (1972) Evaluation of the health aspects of carob bean gum as a food ingredient. Report No. FDABF-GRAS-129. p.1-10.
6. Roukas, T. (1994) Kinetics of ethanol production from carob pods extract by immobilized *Saccharomyces cerevisiae* cells. *Appl Biochem Biotechnol.* 44(1):49-64.
7. Shahkhalili, Y.; Finot, P.A.; Hurrell, R. and Fern, E. (1990) Effects of foods rich in polyphenols on nitrogen excretion in rats. *J Nutr.* 120(4):346-352.
8. Robaislek, E. (1974) Bioavailable calorie assay of guar gum. Unpublished report from WARF Institute, Inc., submitted to the World Health Organization by the Institut Europeen des Industries de la gomme de Caroube.
9. Tsay, L.B. and Whistler, R.L. (1975) Digestibility of galactomannans. Unpublished report submitted to World Health Organization by professor H. Neukom, Chairman of the Technical Committee of the Institut Europeen des Industries de la gomme de Caroube (Cited in FAO/WHO, 1983).
10. Semenza, G. (1975) Report on the possible digestion of the locust bean gum in the stomach and/or in the small intestine in an *in vitro* study. Unpublished report from the Eidgenossische Technische Hochschule Zurich submitted to the World Health Organization by the Institut Europeen des Industries de la gomme de Caroube,
11. Elsenhans, B.; Sufke, U.; Blume, R. and Caspary, W.F. (1980) The influence of carbohydrate gelling agents on rat intestinal transport of monosaccharides and neutral amino acids *in vitro*. *Clin Sci (Lond)* 59(5):373-380.
12. Edwards, C.A.; Blackburn, N.A.; Craigen, L.; Davison, P.; Tomlin, J.; Sugden, K.; Johnson, I.T. and Read, N.W. (1987) Viscosity of food gums determined *in vitro* related to their hypoglycemic actions. *Am J Clin Nutr.* 46(1):72-77.
13. Tsai, A.C. and Peng, B. (1981) Effects of locust bean gum on glucose tolerance, sugar digestion, and gastric motility in rats. *J Nutr.* 111(12):2152-2156.
14. Ershoff, B.H. and Wells, A.F. (1962) Effects of gum guar, locust bean gum and carrageenan on liver cholesterol of cholesterolfed rats. *Proc. Soc Exp. Biol Med* 110:580-582.

15. FAO/WHO (1981) *International Programme on Chemical Safety. Toxicological evaluation of certain food additives and food contaminants. Carob (Locust) bean gum.* WHO Food Additives Series. p.60-69.
16. National Toxicology Program (1981) Carcinogenesis bioassay of locust bean gum (CAS No. 9000-40-2) in F344 rats and B6C3F1 mice (feed study). *National Toxicology Program Technical Report Series.* Report No. 221. p.1-99.
17. Maxwell, W.A. and Newell, G.W. (1972) Study of mutagenic effects of locust bean gum (FDA 71-14). Stanford Research Institute, prepared for the Food and Drug Administration, Menlo Park, California. Report No. NTIS PB-221819. p.1-92.
18. Takada, K.; Toyoda, K.; Shoda, T.; Uneyama, C.; Tamura, T. and Takahashi, M. (1997) [A 13-week subchronic oral toxicity study of carob germ colour in F344 rats]. *Kokuritsu Iyakuhin Shokuhin Eisei Kenkyusho Hokoku* (115):93-98.
19. Cox, G.E.; Baily, D.E. and Morgareidge, K. (1974) Subacute feeding in dogs with a pre-cooked gum blend. Unpublished report from the Food and Drug Laboratories, Inc. submitted to the FAO/WHO Expert Committee.
20. Melnick, R.L.; Huff, J.; Haseman, J.K.; Dieter, M.P.; Grieshaber, C.K.; Wyand, D.S.; Russfield, A.B.; Murthy, A.S.; Fleischman, R.W. and Lilja, H.S. (1983) Chronic effects of agar, guar gum, gum arabic, locust-bean gum, or tara gum in F344 rats and B6C3F1 mice. *Food Chem Toxicol* 21(3):305-311.
21. Brusick, D. (1975) Mutagenic Evaluation of compound: locust bean gum. Unpublished Work.
22. Ashby, J. and Tennant, R.W. (1988) Chemical structure, Salmonella mutagenicity and extent of carcinogenicity as indicators of genotoxic carcinogenesis among 222 chemicals tested in rodents by the U.S. NCI/NTP. *Mutat. Res* 204(1):17-115.
23. Green, S. (1977) Present and future uses of mutagenicity tests for assessment of the safety of food additives. *Journal of Environmental Pathology and Toxicology* 1:49-54.
24. FDRL (1972) Teratologic evaluation of FDA 71-14 (Carob bean (locust) gum). Food and Drug Research Lab., Inc. Report No. NTIS PB221784.
25. Morgareidge, K. (1972) Teratologic evaluation of FDA 71-14 (Carob bean (locust) gum). Food and Drug Research Labs., Inc., Maspeth, New York. Report No. NTIS PB221784. p.1-55.
26. Holbrook, A.A. (1951) The behavior of carob gum in the gastrointestinal tract of man. *Am J Dig. Dis.* 18(1):24-28.
27. Behall, K.M.; Scholfield, D.J.; Lee, K.; Powell, A.S. and Moser, P.B. (1987) Mineral balance in adult men: effect of four refined fibers. *Am J Clin Nutr.* 46(2):307-314.
28. Finn, H. and Haller, H. (1980) [Modification of glucose resorption by bulk materials]. *Z. Gesamte Inn. Med* 35(3):132-136.
29. Kwiterovich, P.O., Jr. (1995) The role of fiber in the treatment of hypercholesterolemia in children and adolescents. *Pediatrics* 96(5 Pt 2):1005-1009.

30. Zavoral, J.H.; Hannan, P.; Fields, D.J.; Hanson, M.N.; Frantz, I.D.; Kuba, K.; Elmer, P. and Jacobs, D.R., Jr. (1983) The hypolipidemic effect of locust bean gum food products in familial hypercholesterolemic adults and children. *Am J Clin Nutr.* 38(2):285-294.
31. Haskell, W.L.; Spiller, G.A.; Jensen, C.D.; Ellis, B.K. and Gates, J.E. (1992) Role of water-soluble dietary fiber in the management of elevated plasma cholesterol in healthy subjects. *Am J Cardiol.* 69(5):433-439.
32. van, d.B., X; Ledent, C. and Mairesse, M. (1992) Rhinitis and asthma caused by occupational exposure to carob bean flour. *J Allergy Clin Immunol.* 90(6 Pt 1):1008-1010.
33. Anonymous (1993) Allergy alert. *Food Safety Notebook* 4:30-32.
34. Scoditti, A.; Peluso, P.; Pezzuto, R.; Giordano, T. and Melica, A. (1996) Asthma to carob bean flour. *Ann. Allergy Asthma Immunol.* 77(1):81.
35. Fiocchi, A.; Restani, P.; Travaini, M.; Decet, E.; Gaiaschi, A.; Bernardo, L. and Riva, E. (1999) Carob is not allergenic in peanut-allergic subjects. *Clin Exp. Allergy* 29(3):402-406.
36. PM USA (2001a) P&T/GC/MS analysis of carob bean extract. Request 20010500. Scan TF131RFB. Unpublished Internal Report.
37. PM USA (2001b) Pyrolysis GC/MS analysis of carob bean extract. Request 20010500. Scan P010500.D. Unpublished Internal Report.
38. INBIFO (2001) In vitro mutagenicity of mainstream smoke condensate of research cigarettes with different amounts of carob bean extract added to the filler: *Salmonella typhimurium* reverse mutation assay. Report No. Institut Fur biologische Forshung Gmbh Report No. P 0500/3341.
39. INBIFO (2001) In vitro cytotoxicity of mainstream smoke fractions of the research cigarette 98.NG.132, 98.NG.133, 98.NG.134, and 98.NG.135: Neutral red uptake assay with mouse embryo BALB/c 3T3 cells. Report No. Project PM Flavor FRR-PM (Institut Fur biologische Forshung Gmbh Report No. P 0500/3337).
40. INBIFO (2000) Chemical analysis of mainstream smoke from cigarettes with 0, 1500, 4500, and 7000 ppm carob bean extract added to the filler. Report No. Project Flavor FRR-PM (Institut Fur biologische Forshung Gmbh Report No. P 0500/3342).
41. CRC (2001) Biological Activity of the Mainstream Smoke of Cigarettes with Different Amounts of Carob Bean Added to the Filler: 90-day Inhalation Study on Rats Projects PM Flavor FRR-PM. Report No. Contract Research Center Report No. P 0500/3329.