

**Evaluation Summary of  
Carob bean  
for Use as a Cigarette Ingredient**

**October 2005**

## **INTRODUCTION**

Carob bean (no CAS assigned) is currently used worldwide at levels below **1,500 ppm** in selected cigarette brands manufactured and/or distributed by Philip Morris International. Carob bean may be applied directly to the tobacco during cigarette manufacturing as a tobacco flavoring material, and as such, may be subject to pyrolysis-type reactions. Carob bean may also be applied to the filter as a flavoring material where it would not be subjected to pyrolysis temperatures. This document summarises our internal studies and current published toxicology information on carob bean abstracted from online toxicity databases.

## **TOXICITY DATA ON UN-BURNED MATERIAL**

### ***Overview***

Carob bean is Generally Recognized As Safe (GRAS) by the U.S. Food and Drug Administration (FDA) for direct food additive in accordance with 21 CFR 184.1343.<sup>1</sup> The Joint WHO/FAO Expert Committee of Food Additives (JECFA) determined that the carob bean gum is safe at historical levels of use in food.<sup>2</sup> Furthermore, carob bean is approved for use by the Council of Europe (CoE No. 120).<sup>3</sup>

Carob bean (sometimes referred to as the powder or flour) is a product derived from milled roasted fruit pod of *Ceratonia siliqua Leguminosae*, a large evergreen tree native to the Middle East region and to the southern part of the Mediterranean. It is extensively cultivated for the seeds and fruit in Cyprus, Greece, Syria, Spain and Italy.<sup>4</sup> The carob tree (also referred to as the locust tree) can reach heights of 33 feet and is well adapted to arid farming conditions.<sup>5</sup> Carob bean (powder or flour) is produced by a continuous process of drying, grinding, and roasting of the pods. The resultant flour has a versatile array of uses in cooking, as well as nutritional and medicinal values. Carob bean powder is almost 50% natural sugar and can be used instead of sugar in most bread and pastry products.<sup>6</sup>

### ***Health hazard data***

A thorough search of the literature on carob bean failed to reveal significant information regarding the potential toxicity of carob bean. Since carob bean gum is used extensively in food manufacturing, most of the relevant published literature data involved studies conducted with carob bean gum. This summary therefore includes data derived from studies with carob bean gum. The relevance of the results of testing with carob bean gum to those that might be produced by carob bean is unknown.<sup>6</sup>

Carob bean gum is chemically defined as a 1,4-galactomannan polymer and biologically characterized as an indigestible polysaccharide hydrocolloid.<sup>7</sup> Carob bean gum moves through the small intestine and arrives in the lower bowel largely undigested.<sup>8-10</sup> Although decreases in serum glucose and cholesterol have been associated with ingestion of large amounts of carob bean gum by experimental animals,<sup>11-14</sup> the research information speaks to a secondary mechanism on these findings – through a reduction in gastric emptying and perhaps peristalsis, thereby delaying or retarding their absorption.<sup>13</sup>

The toxicology studies with carob bean gum further support the inert nature of this molecule. Acute toxicity studies in several species did not show any evidence of oral toxicity at doses up to 10 g/kg.<sup>2,15,16</sup> Subchronic toxicity studies of dose level in rodents up to 10% were unremarkable.<sup>15-17</sup> Similarly, dogs receiving up to 10% (7,500 mg/kg/day) of carob bean gum admixed in the diet for 30 weeks showed no evidence of a compound-related toxicity.<sup>18</sup> Lifetime bioassay studies in rats and mice with up to 5%

(2,500 mg/kg/day for rats and 7,500 mg/kg/day for mice) dietary exposure to carob bean gum did not evoke any meaningful evidence that galactomannan possessed a carcinogenic (or a long-term toxicological) potential.<sup>15,19</sup> Mutagenicity studies were negative<sup>16,20-22</sup> and no chromosomal damage has been demonstrated.<sup>16,22</sup> Teratology studies performed in four species (rats,<sup>23</sup> mice,<sup>23</sup> hamsters<sup>24</sup> and rabbits<sup>24</sup>) gave no suggestion of a teratogenic effect at levels of 910–1,300 mg/kg, although maternal toxicity was evident in rabbits and mice at the highest doses.<sup>23,24</sup>

Carob bean gum has similarly low toxicity in humans. Several clinical studies have failed to show that oral ingestion of carob bean gum has any adverse sequelae.<sup>12,25,26</sup> There is clinical data to support claims of a decrease in serum cholesterol and a blunting of the glucose-tolerance test following oral ingestion of carob bean gum, but whether this is directly related to a carob bean gum effect on gastrointestinal peristalsis or a consequence of the physical nature of the hydrocolloid structure is not known.<sup>12,27-30</sup> The impact of these serum biochemical findings has not been readily embraced by the medical community because carob bean gum is not normally a recommended dietary approach to correcting even marginal increases in plasma cholesterol levels or fasting glucose swings.<sup>6</sup>

The only finding of consequence occurred with carob bean flour, a material derived from the unrefined whole ground seed. There are reports of an allergic response to extensive inhalation exposure to this flour.<sup>31-34</sup> It is not known if this response is a rare occurrence. However, it does follow a pattern that has been noted with inhalation exposure to other proteinaceous materials, including soy protein.<sup>6</sup>

## **TOXICITY DATA ON BURNT MATERIAL**

### ***Combustion studies***

As suggested by the purge and trap studies conducted by PM USA,<sup>35</sup> carob bean applied to tobacco would not be expected to extensively distill at 100°C. At the higher temperatures used in the pyrolysis studies conducted by PM USA, the identified peaks were consistent with chemicals expected from the pyrolytic decomposition of polysaccharides, and suggest that carob bean would be pyrolyzed extensively.<sup>36</sup>

### ***Philip Morris ingredient testing program***

Carob bean was part of a PM USA testing program that was designed to evaluate the potential effects of 333 ingredients added to typical commercial blended test cigarettes on selected biological and chemical endpoints.<sup>37-40</sup> Three pairs of test cigarettes were produced, each containing different groups of ingredients. Carob bean was added to one pair at target levels of 1,165 and 1,748 ppm. No significant effects were noted in cytotoxicity, mutagenic studies or in respiratory tract endpoints in 90-day rat inhalation studies. In addition, smoke chemistry studies from cigarettes containing a mixture of flavors including carob bean did not significantly alter the smoke chemistry profile compared to control cigarettes. Based on the results of these studies, the authors concluded that these ingredients (including carob bean) added to tobacco do not add significantly to the overall toxicity of cigarettes.

## **CONCLUSION**

Cigarette smoking causes lung cancer, heart disease, emphysema and other serious diseases in smokers. Smokers are far more likely to develop serious diseases, like lung cancer, than non-smokers. There is no "safe" cigarette. Government health warnings about smoking apply to all cigarettes, regardless of the ingredients added, including those containing only tobacco and paper.

Currently, information is only available for tests utilizing carob bean at levels below the current use level. Studies are ongoing to address the use of carob bean at higher tobacco application levels. While the current use level exceeds the levels that have been tested, published studies show there is no meaningful difference in the composition or toxicity of smoke from cigarettes with added ingredients (including carob bean) compared to the smoke from cigarettes without added ingredients.<sup>37-44</sup> It is our scientific judgement, based on the best available data, that carob bean used in our cigarettes does not increase the overall toxicity of cigarette smoke.

## References

1. Food and Drug Administration (1974) Part 121 - Food Additives. Subpart B - Exemption of certain food additives from the requirement of tolerances. Locust (carob) bean gum. Report No. Federal Register 39. p.34181-34184.
2. FAO/WHO (1981) *International Programme on Chemical Safety. Toxicological evaluation of certain food additives and food contaminants. Carob (Locust) bean gum.* WHO Food Additives Series. p.60-69.
3. CoE (1981) *Flavouring Substances and Natural Sources of Flavourings.* Council of Europe Publishing, F-67006 Strausbourg Cedex, France. p.59-113.
4. LSRO (1972) Evaluation of the health aspects of carob bean gum as a food ingredient. Report No. FDABF-GRAS-129. p.1-10.
5. Roukas, T. (1994) Kinetics of ethanol production from carob pods extract by immobilized *Saccharomyces cerevisiae* cells. *Appl Biochem Biotechnol.* 44(1):49-64.
6. Burdock, G.A. (2000) Safety Assessment of Carob Bean Gum as a Food Ingredient. Burdock Group, Vero Beach, FL. Unpublished Review.
7. Shahkhalili, Y.; Finot, P.A.; Hurrell, R. and Fern, E. (1990) Effects of foods rich in polyphenols on nitrogen excretion in rats. *J Nutr.* 120(4):346-352.
8. Robaislek, E. (1974) Bioavailable calorie assay of guar gum. Unpublished report from WARF Institute, Inc., submitted to the World Health Organization by the Institut Europeen des Industries de la gomme de Caroube.
9. Tsay, L.B. and Whistler, R.L. (1975) Digestibility of galactomannans. Unpublished report submitted to World Health Organization by professor H. Neukom, Chairman of the Technical Committee of the Institut Europeen des Industries de la gomme de Caroube.
10. Semenza, G. (1975) Report on the possible digestion of the locust bean gum in the stomach and/or in the small intestine in an *in vitro* study. Unpublished report from the Eidgenossische Technische Hochschule Zurich submitted to the World Health Organization by the Institut Europeen des Industries de la gomme de Caroube,
11. Elsenhans, B.; Sufke, U.; Blume, R. and Caspary, W.F. (1980) The influence of carbohydrate gelling agents on rat intestinal transport of monosaccharides and neutral amino acids *in vitro.* *Clin Sci (Lond)* 59(5):373-380.
12. Edwards, C.A.; Blackburn, N.A.; Craigen, L.; Davison, P.; Tomlin, J.; Sugden, K.; Johnson, I.T. and Read, N.W. (1987) Viscosity of food gums determined *in vitro* related to their hypoglycemic actions. *Am J Clin Nutr.* 46(1):72-77.
13. Tsai, A.C. and Peng, B. (1981) Effects of locust bean gum on glucose tolerance, sugar digestion, and gastric motility in rats. *J Nutr.* 111(12):2152-2156.
14. Ershoff, B.H. and Wells, A.F. (1962) Effects of gum guar, locust bean gum and carrageenan on liver cholesterol of cholesterolfed rats. *Proc. Soc Exp. Biol Med* 110:580-582.

15. National Toxicology Program (1981) Carcinogenesis bioassay of locust bean gum (CAS No. 9000-40-2) in F344 rats and B6C3F1 mice (feed study). *National Toxicology Program Technical Report Series*. Report No. 221. p.1-99.
16. Maxwell, W.A. and Newell, G.W. (1972) Study of mutagenic effects of locust bean gum (FDA 71-14). Stanford Research Institute, prepared for the Food and Drug Administration, Menlo Park, California. Report No. NTIS PB-221819. p.1-92.
17. Takada, K.; Toyoda, K.; Shoda, T.; Uneyama, C.; Tamura, T. and Takahashi, M. (1997) [A 13-week subchronic oral toxicity study of carob germ colour in F344 rats]. *Kokuritsu Iyakuhin Shokuhin Eisei Kenkyusho Hokoku* (115):93-98.
18. Cox, G.E.; Baily, D.E. and Morgareidge, K. (1974) Subacute feeding in dogs with a pre-cooked gum blend. Unpublished report from the Food and Drug Laboratories, Inc. submitted to the FAO/WHO Expert Committee.
19. Melnick, R.L.; Huff, J.; Haseman, J.K.; Dieter, M.P.; Grieshaber, C.K.; Wyand, D.S.; Russfield, A.B.; Murthy, A.S.; Fleischman, R.W. and Lilja, H.S. (1983) Chronic effects of agar, guar gum, gum arabic, locust-bean gum, or tara gum in F344 rats and B6C3F1 mice. *Food Chem Toxicol* 21(3):305-311.
20. Brusick, D. (1975) Mutagenic Evaluation of compound: locust bean gum. Unpublished Work.
21. Ashby, J. and Tennant, R.W. (1988) Chemical structure, Salmonella mutagenicity and extent of carcinogenicity as indicators of genotoxic carcinogenesis among 222 chemicals tested in rodents by the U.S. NCI/NTP. *Mutat. Res* 204(1):17-115.
22. Green, S. (1977) Present and future uses of mutagenicity tests for assessment of the safety of food additives. *Journal of Environmental Pathology and Toxicology* 1:49-54.
23. FDRL (1972) Teratologic evaluation of FDA 71-14 (Carob bean (locust) gum). Food and Drug Research Lab., Inc. Report No. NTIS PB221784.
24. Morgareidge, K. (1972) Teratologic evaluation of FDA 71-14 (Carob bean (locust) gum). Food and Drug Research Labs., Inc., Maspeth, New York. Report No. NTIS PB221784. p.1-55.
25. Holbrook, A.A. (1951) The behavior of carob gum in the gastrointestinal tract of man. *Am J Dig. Dis.* 18(1):24-28.
26. Behall, K.M.; Scholfield, D.J.; Lee, K.; Powell, A.S. and Moser, P.B. (1987) Mineral balance in adult men: effect of four refined fibers. *Am J Clin Nutr.* 46(2):307-314.
27. Finn, H. and Haller, H. (1980) [Modification of glucose resorption by bulk materials]. *Z. Gesamte Inn. Med* 35(3):132-136.
28. Kwiterovich, P.O., Jr. (1995) The role of fiber in the treatment of hypercholesterolemia in children and adolescents. *Pediatrics* 96(5 Pt 2):1005-1009.
29. Zavoral, J.H.; Hannan, P.; Fields, D.J.; Hanson, M.N.; Frantz, I.D.; Kuba, K.; Elmer, P. and Jacobs, D.R., Jr. (1983) The hypolipidemic effect of locust bean gum food products in familial hypercholesterolemic adults and children. *Am J Clin Nutr.* 38(2):285-294.

30. Haskell, W.L.; Spiller, G.A.; Jensen, C.D.; Ellis, B.K. and Gates, J.E. (1992) Role of water-soluble dietary fiber in the management of elevated plasma cholesterol in healthy subjects. *Am J Cardiol.* 69(5):433-439.
31. van, d.B., X; Ledent, C. and Mairesse, M. (1992) Rhinitis and asthma caused by occupational exposure to carob bean flour. *J Allergy Clin Immunol.* 90(6 Pt 1):1008-1010.
32. Anonymous (1993) Allergy alert. *Food Safety Notebook* 4:30-32.
33. Scoditti, A.; Peluso, P.; Pezzuto, R.; Giordano, T. and Melica, A. (1996) Asthma to carob bean flour. *Ann. Allergy Asthma Immunol.* 77(1):81.
34. Fiocchi, A.; Restani, P.; Travaini, M.; Decet, E.; Gaiaschi, A.; Bernardo, L. and Riva, E. (1999) Carob is not allergenic in peanut-allergic subjects. *Clin Exp. Allergy* 29(3):402-406.
35. PM USA (2001a) P&T/GC/MS analysis of carob bean. Request 20010380. Scan TF061RFB. Unpublished Internal Report.
36. PM USA (2001b) Pyrolysis GC/MS analysis of carob bean. Request 20010380. Scan 01DS231.D. Unpublished Internal Report.
37. Carmines, E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 1: Cigarette design, testing approach, and review of results. *Food and Chemical Toxicology* 40:77-91.
38. Roemer, E.; Tewes, F.J.; Meisgen, T.J.; Veltel, D. and Carmines, E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 3: *In vitro* genotoxicity and cytotoxicity. *Food and Chemical Toxicology* 40:105-111.
39. Rustemeier, K.; Stabbert, R.; Haussmann, H.J.; Roemer, E. and Carmines E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 2: Chemical composition of mainstream smoke. *Food and Chemical Toxicology* 40:93-104.
40. Vanscheeuwijck, P.M.; Teredesai, A.; Terpstra, P.M.; Verbeek, J.; Kuhl, P.; Gerstenberg, B.; Gebel, S. and Carmines E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 4: Subchronic inhalation toxicity. *Food and Chemical Toxicology* 40:113-131.
41. Gaworski, C.L.; Dozier, M.M.; Heck, J.D.; Gerhart, J.M.; Rajendran, N.; David, R.M.; Brennecke, L.H. and Morrissey, R. (1998) Toxicologic evaluation of flavor ingredients added to cigarette tobacco: 13 week inhalation exposures in rats. *Inhal. Toxicol.* 10:357-381.
42. Gaworski, C.L.; Heck, J.D.; Bennett, M.B. and Wenk, M.L. (1999) Toxicologic evaluation of flavor ingredients added to cigarette tobacco: skin painting bioassay of cigarette smoke condensate in SENCAR mice. *Toxicology* 139(1-2):1-17.
43. Doull, J.; Frawley, J.P.; George, W.J.; Loomis, T.A.; Squire, R.A. and Taylor, S.L. (1994) A safety assessment of ingredients added to tobacco in the manufacturing of cigarettes. Covington and Burling, Washington, D.C.
44. Doull, J.; Frawley, J.P.; George, W.J.; Loomis, T.A.; Squire, R.A. and Taylor, S.L. (1998) A safety assessment of ingredients added to tobacco in the manufacturing of cigarettes. Covington and Burling, Washington, D.C.