

**Evaluation Summary of
Citric acid
for Use as a Cigarette Ingredient**

October 2005

INTRODUCTION

Citric acid (CAS # 77-92-9) is currently used worldwide at levels below **8,000 ppm** in selected cigarette brands manufactured and/or distributed by Philip Morris International. Citric acid is a natural component of tobacco.¹⁷ Citric acid is also used in tobacco manufacturing as a flavoring agent, and may also be found as a minor component of adhesives and papers. As such, citric acid may be subject to pyrolysis-type reactions. This document summarises our internal studies and current published toxicology information on citric acid abstracted from online toxicity databases.

TOXICITY DATA ON UN-BURNED MATERIAL

Overview

Citric acid occurs in many foods and has been used widely in the food industry for more than 125 years as an acidulent, sequestering agent, synergistic antioxidant, dispersing agent, flavor enhancer, and water-conditioning agent.¹ Both the U.S. Food and Drug Administration (21 CFR §184.1033) and the Flavor and Extract Manufacturers Association (FEMA No. 2306)² have designated citric acid as generally recognized as safe (GRAS) for use in food.

Health hazard data

Citric acid has a relatively low order of acute toxicity in rodents.³⁻⁷ Long-term feeding studies in rats at concentrations up to 5% of the diet indicated slightly reduced growth, but no tissue abnormalities nor evidence of carcinogenicity.^{4,7-11} Inhalation of high concentrations of citric acid aerosols has been reported to cause coughing in humans.^{7,12} Citric acid has not been reported to be a reproductive or developmental toxicant.^{3,7,8,11,13} Citric acid was not mutagenic in the Salmonella bacteria assay.^{7,14-16} Citric acid did not induce chromosomal damage in cultured human or hamster cells.^{7,14} In humans, direct application of citric acid crystals to the inside of the mouth has been reported to cause ulcers while skin irritation has been reported among occupationally exposed waiters and bakers.⁷

TOXICITY DATA ON BURNT MATERIAL

Combustion studies

As suggested by the purge and trap studies conducted by PM USA, citric acid applied to tobacco would not be expected to significantly distill at 100°C.¹⁸ At the higher temperatures used in the PM USA pyrolysis studies; the two largest peaks were identified as carbon dioxide and citraconic anhydride. These peaks in addition to the smaller acetone peak suggest that citric acid would be pyrolyzed and would not be delivered to the smoke intact.¹⁹

Philip Morris ingredient testing program

Citric acid was a part of the PM USA ingredient testing program that was designed to evaluate the potential effects of ingredients added to typical commercial blended test cigarettes on selected biological and chemical endpoints. Citric acid was added to test cigarette tobacco at target concentrations of 5,000, 18,000 or 42,000 ppm above the natural background found in tobacco. Application of target levels of 42,000 ppm or 18,000 ppm citric acid to test cigarette tobacco increased the mainstream concentrations of some smoke constituents including hydrogen cyanide, formaldehyde, acrolein and polycyclic aromatic

hydrocarbons. Application of a target level of 5,000 ppm citric acid did not produce these increases. Although some statistically significant differences were noted between control cigarettes and cigarettes with high target levels of citric acid, the increased smoke constituent concentrations were considered to be within normal variation of cigarette products.²⁰ The mutagenic response of *Salmonella* bacteria exposed to mainstream smoke condensate samples obtained from cigarettes with citric acid was not significantly different from condensate from control cigarettes.²¹ Similarly, the cytotoxic response of mouse embryo BALB/c 3T3 cells treated with mainstream smoke condensate preparations was not altered by citric acid addition.²² The biological effects of inhaling smoke from cigarettes with citric acid was assessed in Sprague-Dawley rats exposed nose-only to smoke for 6 hrs/day, 7 days/week for 13 weeks. The results of the smoke inhalation studies indicated that citric acid addition to cigarette tobacco at the levels tested did not discernibly alter the biological effects normally associated with cigarette smoke exposure in rodents.²³

CONCLUSION

Cigarette smoking causes lung cancer, heart disease, emphysema and other serious diseases in smokers. Smokers are far more likely to develop serious diseases, like lung cancer, than non-smokers. There is no "safe" cigarette. Government health warnings about smoking apply to all cigarettes, regardless of the ingredients added, including those containing only tobacco and paper.

The results of this evaluation of citric acid involving a review of published information and internal studies show there is no meaningful difference in the composition or toxicity of smoke from cigarettes with added ingredients (including citric acid) compared to the smoke from cigarettes without added ingredients.²⁰⁻²⁴ It is our scientific judgement, based on the best available data, that citric acid used in our cigarettes does not increase the overall toxicity of cigarette smoke.

References

1. Joint FAO/WHO Expert Committee on Food Additives (1974) *Toxicological evaluation of some food additives including anticaking agents, antimicrobials, antioxidants, emulsifiers and thickening agents*. World Health Organization, Geneva.
2. Hall, R.L. and Oser, B.L. (1965) Recent Progress in the Consideration of Flavoring Ingredients Under the Food Additives Amendment. III GRAS Substances. *Food Technology* 19:151-197.
3. Zhang, Y. and Cheng Z. (1987) Teratogenic effect of citric acid in rats. *Shanghai Yike Daxue Xuebao*. 14:195-198.
4. Yokotani, H.; Usui, T.; Nakaguchi, T.; Kanabayashi, T.; Tanda, M. and Aramaki, Y. (1971) Acute And Subacute Toxicological Studies Of TAKEDA-Citric Acid In Mice And Rats. *Takeda Kenkyusho Ho* :
5. Gruber, C.M. and Halbeisen, W.A. (1948) A study on the comparative toxic effects of citric acid and its sodium salts. *Journal of Pharmacology and Experimental Therapy* 94:65-67.
6. Oelkers, H.A. (1965) Contribution to the pharmacology of piperazine. *Arztl. Forsch* 19(12):625-630.
7. BIBRA (1993) Citric acid and its common salts.
8. Bonting, S.L. and Jansen, B.C.P. (1956) The effect of a prolonged intake of phosphoric acid and citric acid in rats. *Voeding* (17):137-148.
9. Fukushima, S.; Thamavit, W.; Kurata, Y. and Ito, N. (1986) Sodium citrate: a promoter of bladder carcinogenesis. *Jpn J Cancer Res* 77(1):1-4.
10. Wright, E. and Hughes, R.E. (1976) Some effects of dietary citric acid in small animals. *Food Cosmet Toxicol* 14(6):561-564.
11. Wright, E. and Hughes, R.E. (1976) Influence of dietary citric acid supplement on reproduction and survival time of mice and rats. *Nutrition Reports International*. 13(6):563-566.
12. Bickerman, H.A. and Barach, A.L. (1954) The experimental production of cough in human subjects induced by citric acid aerosols. Preliminary studies on the evaluation of antitussive agents. *American Journal of Medical Science*. (228):156-163.
13. Gomez, M.; Domingo, J.L. and Llobet, J.M. (1991) Developmental toxicity evaluation of oral aluminum in rats: influence of citrate. *Neurotoxicol. Teratol* :
14. Ishidate, M., Jr.; Sofuni, T.; Yoshikawa, K.; Hayashi, M.; Nohmi, T.; Sawada, M. and Matsuoka, A. (1984) Primary mutagenicity screening of food additives currently used in Japan. *Food Chem Toxicol* 22(8):623-636.
15. Al Ani, F.Y. and Al Lami, S.K. (1988) Absence of mutagenic activity of acidity regulators in the Ames Salmonella/microsome test. *Mutat. Res* 206(4):467-470.
16. Bala, S. and Grover, I.S. (1989) Antimutagenicity of some citrus fruits in Salmonella typhimurium. *Mutat. Res* 222(3):141-148.

17. Leffingwell, J.C. (1999) Leaf chemistry: Basic chemical constituents of tobacco leaf and differences among tobacco types. In *Tobacco Production, Chemistry and Technology*. (D.L.Davis and M.T.Nielsen, Eds.). Blackwell Sciences Ltd., Oxford. p.265-284.
18. PM USA (2001) P&T/GC/MS Analysis of Citric Acid. Request 20010601. Scan TG181HCB.D. Unpublished Internal Report.
19. Philip Morris USA Internal Report (2001) Pyrolysis GC/MS analysis of citric acid. Request 20010601 File P010601A.D.
20. INBIFO (1999) Institut Fur biologische Forshung Gmbh. Chemical analysis of mainstream smoke from cigarettes 98.FB.107, 98.FB.109, 98.FB.110, and 98.FB.111: Project JUICE. Report No. P 0500/3316.
21. INBIFO (2001) Institut Fur biologische Forshung Gmbh. In Vitro Mutagenicity of Mainstream Smoke Condensate of the Research Cigarettes 98.FB.107, 98.FB.109, 98.FB.110, 98.FB.111, 98.FB.113, 98.FB.114, and 98.FB.116 Salmonella Typhimurium Reverse Mutation Assay: Projects MILK and JUICE. Report No. P 0500/3303.
22. INBIFO (1999) Institut Fur biologische Forshung Gmbh. In Vitro Cytotoxicity of mainstream smoke fractions of the research cigarettes 98.FB.107, 98.FB.109, 98.FB.110, 98.FB.111, 98.FB.113, 98.FB.114, and 98.FB.116, Neutral red uptake assay with mouse embryo BALB/c 3T3 cells: Projects MILK and JUICE. Report No. P 0500/3298.
23. CRC (2000) Contract Research Center. Biological Activity of the Mainstream Smoke of the Test Cigarettes 98.FB.109, 98.FB.110, 98.FB.111, 98.FB.113, 98.FB.114, 98.FB.116, and the Control Cigarette 98.FB.107 90-Day Inhalation Study on Rats Projects MILK and JUICE. Report No. B3299 P 0500/3299.
24. Baker, R.R.; Massey, E.D. and Smith, G. (2004) An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food and Chemical Toxicology* 42S(Supplement 1):S53-S83.