

**Evaluation Summary of
Cocoa
for Use as a Cigarette Ingredient**

October 2005

INTRODUCTION

Cocoa (CAS # 8002-31-1) is currently used worldwide at levels below **9,000 ppm** in selected cigarette brands manufactured and/or distributed by Philip Morris International. As such, cocoa may be subject to pyrolysis-type reactions when smoked. Cocoa may also be applied to the filter as a flavoring material where it would not be subjected to pyrolysis temperatures. This document summarises our internal studies and current published toxicology information on cocoa abstracted from online toxicity databases.

TOXICITY DATA ON UN-BURNED MATERIAL

Overview

Cocoa is the basis of chocolate flavoring and is the confection of choice in the United States, representing nearly half of the total candy consumption. Cocoa extract is also consumed at high levels, placing it among the top 16% of food ingredients when ranked according to consumption.^{1,2} Cocoa powder, cocoa butter, and cocoa liquor are all products of the same fruit from the cacao tree, a member of the family *Sterculiaceae*, most often *Theobroma cacao* L. and rarely, *T. pentagona* or *T. sphaerocarpa*.³ Cocoa is regulated as a food by the U.S. Food and Drug Administration (21 CFR § 163.110 thru 21 CFR § 163.155).

Cocoa butter is unlikely to be toxic at average human consumption levels due to the low reported toxicities of the individual fatty acids contained in the butter.⁴⁻⁶ The high content of long chain fatty acids (oleic and stearic acids) in cocoa butter have been linked to their hypocholesterolemic effects in study animals.^{7,8}

Health hazard data

Animal studies of cocoa powder toxicity have related its effects to the amount of theobromine contained in the cocoa.^{3,6,9-11} Theobromine is considered the most relevant pharmacologically active agent contained in cocoa powder.^{11,12} Animal studies have indicated that theobromine is toxic to Sertoli cells of the testes in rats.^{13,14} In comparison to other commonly consumed methylxanthines, caffeine and theophylline, theobromine has a low potency in regards to its effects on cytosolic calcium levels, adenosine receptor antagonism and inhibition of phosphodiesterases.^{11,12} Theobromine is not considered to be a stimulant of the central nervous system as are caffeine and theophylline.^{11,15-22}

Results of reproductive studies in rats and rabbits indicate that high levels of theobromine are required in order to reduce the reproductive viability in these animals. The amounts of cocoa required to reach these same levels of theobromine in humans far surpasses its average daily consumption.²³⁻²⁸ A two year toxicity study found no evidence of carcinogenesis in rats fed diets of up to 5.0% cocoa powder.²⁹

TOXICITY DATA ON BURNT MATERIAL

Combustion studies

PM USA purge and trap studies suggest that cocoa would not distill in front of the burning cone at low temperatures.³⁰ At the higher temperatures used in pyrolysis studies conducted by PM USA, cocoa would be expected to pyrolyze extensively. The major peak identified in the pyrolysis study was theobromine which suggests that the theobromine would be expected to transfer to smoke intact.³¹ Theobromine is a

vasoactive amine³² and the expected exposure level of theobromine from cigarette smoke is considered to be too low to produce a clinically relevant blood level of theobromine.

Philip Morris ingredient testing program

Cocoa was part of a testing program that was designed to evaluate the potential effects of 333 ingredients added to typical commercial blended test cigarettes on selected biological and chemical endpoints. Three pairs of test cigarettes were produced, each containing different groups of ingredients. Cocoa was added to one pair of test cigarettes at target levels of 6470 ppm and 9705 ppm. No significant effects were noted in cytotoxicity, mutagenic studies or in respiratory tract endpoints in 90-day rat inhalation studies. In addition, smoke chemistry studies from cigarettes containing a mixture of flavors (including cocoa) did not significantly alter the smoke chemistry profile compared to control cigarettes. Based on the results of these studies, the authors concluded that these ingredients (including cocoa) added to tobacco do not add significantly to the overall toxicity of cigarettes.³³⁻³⁶

CONCLUSION

Cigarette smoking causes lung cancer, heart disease, emphysema and other serious diseases in smokers. Smokers are far more likely to develop serious diseases, like lung cancer, than non-smokers. There is no "safe" cigarette. Government health warnings about smoking apply to all cigarettes, regardless of the ingredients added, including those containing only tobacco and paper.

Currently, information is only available for tests utilizing cocoa in a mixture of ingredients applied to cigarette tobacco. Studies are ongoing to address the use of cocoa as a single ingredient. Published studies show there is no meaningful difference in the composition or toxicity of smoke from cigarettes with added ingredients (including cocoa) compared to the smoke from cigarettes without added ingredients.³³⁻⁴² It is our scientific judgement, based on the best available data, that cocoa used in our cigarettes does not increase the overall toxicity of cigarette smoke.

References

1. USDA (1997) Foods Commonly Eaten in the United States. Quantities consumed per Eating Occasion and in a Day, 1989-91. United States Department of Agriculture Report No. NTIS PB98-111719.
2. USDA-ERS (1996) Food Consumption, Prices and Expenditures, 1996. Annual Data, 1970-94. United States Department of Agriculture, Economic Research Service, Statistical Bulletin, Number 928 Report No. NTIS PB96-165584.
3. IARC (1991) *Coffee, tea, mate, methylxanthines and methylglyoxyl*. International Agency for Research on Cancer Monograph on the Evaluation of Carcinogenic Risks to Humans. World Health Organization. p.513.
4. Budavari, S.; O'Neil, M.J.; Smith, A. and Heckelman, P.E. (1989) *The Merck Index. An Encyclopedia of Chemicals, drugs and Biologicals*. Eleventh edition Edition. Merck & Co., Inc., Rahway, N.J.
5. Clydesdale, F.M.Ed. (1997) Cocoa Extract. Food Additives. In *Toxicology, regulation and properties*. CRC Press, Boca Raton, FL. CD-ROM. p.CD-ROM.
6. RTECS (1999) Registry of Toxic Effects of Chemical Substances. A database supplied by the National Library of Medicine, Bethesda, MD.
7. Kris-Etherton, P.M.; Derr, J.; Mitchell, D.C.; Mustad, V.A.; Russell, M.E.; McDonnell, E.T.; Salabsky, D. and Pearson, T.A. (1993) The role of fatty acid saturation on plasma lipids, lipoproteins, and apolipoproteins: I. Effects of whole food diets high in cocoa butter, olive oil, soybean oil, dairy butter, and milk chocolate on the plasma lipids of young men. *Metabolism* 42(1):121-129.
8. Kris-Etherton, P.M.; Derr, J.A.; Mustad, V.A.; Seligson, F.H. and Pearson, T.A. (1994) Effects of a milk chocolate bar per day substituted for a high-carbohydrate snack in young men on an NCEP/AHA Step 1 Diet. *Am J Clin Nutr.* 60(6 Suppl):1037S-1042S.
9. Kondo, K.; Hirano, R.; Matsumoto, A.; Igarashi, O. and Itakura, H. (1996) Inhibition of LDL oxidation by cocoa. *Lancet* 348(9040):1514.
10. Shively, C.A. and Tarka, S.M., Jr. (1984) Methylxanthine composition and consumption patterns of cocoa and chocolate products. *Prog. Clin Biol Res* 158:149-178.
11. Tarka, S.M., Jr. (1982) The toxicology of cocoa and methylxanthines: a review of the literature. *Crit Rev Toxicol* 9(4):275-312.
12. Rall, T.W. (1991) Drugs used in the treatment of asthma. In *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. eighth edition Edition. (A. G. Gilman, T. W. Rall, A. S. Nies and P. e. Taylor, Eds.). Pergamon Press, New York, NY. p.618-630.
13. Wang, Y.; Waller, D.P.; Hikim, A.P. and Russell, L.D. (1992) Reproductive toxicity of theobromine and cocoa extract in male rats. *Reprod. Toxicol* 6(4):347-353.
14. Wang, Y. and Waller, D.P. (1994) Theobromine toxicity on Sertoli cells and comparison with cocoa extract in male rats. *Toxicol Lett* 70(2):155-164.

15. Beer, B.; Chasin, M.; Clody, D.E.; Vogel, J.R. and Horovitz, Z.P. (1972) Cyclic monophosphate phosphodiesterase in brain: effect on anxiety. *Science* 176:428-430.
16. Benton, D. (1999) Chocolate craving: biological or psychological phenomenon? In *Chocolate & Cocoa, Health & Nutrition*. (I. e. Knight, Ed.). Blackwell Science, Inc., Malden, MA. p.256-278.
17. Carney, J.M.; Holloway, F.A. and Modrow, H.E. (1985) Discriminative stimulus properties of methylxanthines and their metabolites in rats. *Life Sci* 36(10):913-920.
18. Herz, A.e.al. (1968) Vergleichende Untersuchungen über zentrale Wirkungen von Xanthinderivaen in Hinblick auf deren Stoffwechsel und Verteilung im Organismus. *Naunyn Schmiedebergs Arch. Pharmakol. Exp. Pathol.* 261:1123-1132 (cited in Benton, D.(1999).
19. Holtzman, S.G. (1986) Discriminative stimulus properties of caffeine in the rat: noradrenergic mediation. *J Pharmacol Exp. Ther* 239(3):706-714.
20. Katims, J.J.; Annau, Z. and Snyder, S.H. (1983) Interactions in the behavioral effects of methylxanthines and adenosine derivatives. *J Pharmacol Exp. Ther* 227(1):167-173.
21. Sprugel, W.; Mitznegg, P. and Heim, F. (1977) The influence of caffeine and theobromine on locomotive activity and the brain cGMP/cAMP ratio in white mice. *Biochem Pharmacol* 26(18):1723-1724.
22. Starvic, B. (1988) Methylxanthines: toxicity to humans. 3. Theobromines, paraxanthines and the combined effects of methylxanthines. *Food and Chemical Toxicology* 26:725-733.
23. Hostetler, K.A.; Morrissey, R.B.; Tarka, S.M., Jr.; Apgar, J.L. and Shively, C.A. (1990) Three-generation reproductive study of cocoa powder in rats. *Food Chem Toxicol* 28(7):483-490.
24. Lamb, J.; Gulati, D.; Choudhury, H.; Chambers, R.; Poonacha, K. and Sabharwal, P. (1997) Reproductive toxicology; Theobromine. *Environmental Health Perspectives* 105(suppl 1):353-354.
25. Shively, C.A.; White, D.M.; Blauch, J.L. and Tarka, S.M., Jr. (1984) Dominant lethal testing of theobromine in rats. *Toxicol Lett* 20(3):325-329.
26. Tarka, S.M., Jr.; Applebaum, R.S. and Borzelleca, J.F. (1986) Evaluation of the perinatal, postnatal and teratogenic effects of cocoa powder and theobromine in Sprague-Dawley/CD rats. *Food Chem Toxicol* 24(5):375-382.
27. Tarka, S.M., Jr.; Applebaum, R.S. and Borzelleca, J.F. (1986) Evaluation of the teratogenic potential of cocoa powder and theobromine in New Zealand White rabbits. *Food Chem Toxicol* 24(5):363-374.
28. Tesh, J.M.; Lambert, E.P.; Elias, P.S. and Gottschalk, H.M. (1982) Cocoa beans: effects upon reproductive function, growth and development in the rat. *Teratology* 26(3):21a.
29. Tarka, S.M., Jr.; Morrissey, R.B.; Apgar, J.L.; Hostetler, K.A. and Shively, C.A. (1991) Chronic toxicity/carcinogenicity studies of cocoa powder in rats. *Food Chem Toxicol* 29(1):7-19.
30. PM USA (2001) P&T/GC/MS Analysis of Cocoa. Scan TH221HCD. Request 20010726. Unpublished Internal Report.

31. PM USA (2001) Pyrolysis GC/MS Analysis of Cocoa. Scan P010726A.D. Unpublished Internal Report.
32. Scharff, L. and Marcus, D.A. (1999) Chocolate and Headache: Is there a relationship? In *Chocolate & Cocoa, Health & Nutrition*. (I. e. Knight, Ed.). Blackwell Science, Inc., Malden, MA. p.229-239.
33. Carmines, E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 1: Cigarette design, testing approach, and review of results. *Food and Chemical Toxicology* 40:77-91.
34. Roemer, E.; Tewes, F.J.; Meisgen, T.J. and Veltel, D.a.C.E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 3: *In vitro* genotoxicity and cytotoxicity. *Food and Chemical Toxicology* 40:105-111.
35. Rustemeier, K.; Stabbert, R.; Haussmann, H.J. and Roemer, E.a.C.E.L. (2002) Evaluation of the potential effects of ingredients added to cigarette. Part 2: Chemical composition of mainstream smoke. *Food and Chemical Toxicology* 40:93-104.
36. Vanscheeuwijck, P.M.; Terpstra, P.M.; Teredesai, A.; Verbeek, J.; Kuhl, P.; Gerstenberg, B.; Gebel, S. and and Carmines, E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 4: Subchronic inhalation toxicity. *Food and Chemical Toxicology* 40:113-131.
37. Doull, J.; Frawley, J.P.; George, W.J.; Loomis, T.A.; Squire, R.A. and and Taylor, S.L. (1994) A safety assessment of ingredients added to tobacco in the manufacturing of cigarettes. Covington and Burling, Washington, D.C.
38. Doull, J.; Frawley, J.P.; George, W.J.; Loomis, T.A.; Squire, R.A. and and Taylor, S.L. (1998) A safety assessment of ingredients added to tobacco in the manufacturing of cigarettes. Covington and Burling, Washington, D.C.
39. Roemer, E. and Hackenberg, U. (1990) Mouse skin bioassay of smoke condensates from cigarettes containing different levels of cocoa. *Food Addit. Contam* 7(4):563-569.
40. Gaworski, C.L.; Dozier, M.M.; Heck, J.D.; Gerhart, J.M.; Rajendran, N.; David, R.M.; Brennecke, L.H. and Morrissey, R. (1998) Toxicologic evaluation of flavor ingredients added to cigarette tobacco: 13-week inhalation exposures in rats. *Inhalation Toxicology* 10:357-381.
41. Gaworski, C.L.; Heck, J.D.; Bennett, M.B. and Wenk, M.L. (1999) Toxicologic evaluation of flavor ingredients added cigarette tobacco: Skin painting bioassay of cigarette smoke condensate in SENCAR mice. *Toxicology* 139:1-17.
42. Baker, R.R.; Massey, E.D. and Smith, G. (2004) An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food and Chemical Toxicology* 42S:S53-S83.