

**Evaluation Summary of  
Ethyl acetate  
for Use as a Cigarette Ingredient**

**October 2005**

## **INTRODUCTION**

Currently, ethyl acetate (CAS # 141-78-6) is used worldwide at levels below **100 ppm** in selected cigarette brands manufactured and/or distributed by Philip Morris International. Ethyl acetate is applied directly to the tobacco as an additive, flavoring, flavoring agent, or solvent, and as such, ethyl acetate may be subject to pyrolysis-type reactions when smoked. Ethyl acetate may also be applied to the filter as a flavoring material where it would not be subjected to pyrolysis temperatures. This document summarises our internal studies and current published toxicology information on ethyl acetate abstracted from online toxicity databases.

## **TOXICITY DATA ON UN-BURNED MATERIAL**

### ***Overview***

Ethyl acetate has been approved for addition to food by the U.S. Food and Drug Administration (FDA 21 CFR § 182.60), the Flavour and Extract Manufacturers Association (FEMA No. 2414),<sup>1</sup> the Joint FAO/WHO Expert Committee on Food Additives (JECFA)<sup>2</sup> and the Council of Europe (CoE No. 191).<sup>3</sup> Ethyl acetate has been added to foods since at least 1920, and has been reported to occur in nature.<sup>4-11</sup> It is also used as solvent in nail polish, nail polish removers, basecoats and other manicuring preparations.<sup>12-14</sup>

### ***Health hazard data***

No mutagenic activity of ethyl acetate was detected by Ames test using *Salmonella typhimurium* strains TA92, TA1535, TA1537, TA94, TA98 and for some samples, TA2637, with and without S9 liver microsome fraction.<sup>15-18</sup> Ethyl acetate was nonmutagenic when tested by Rec-assay.<sup>19</sup> However, weakly positive results were obtained in a chromosomal aberration test using a Chinese hamster fibroblasts cell line<sup>16</sup> and in a mitotic aneuploidy test using *S. cerevisiae*.<sup>20,21</sup> Chromosomal aberration positives were designated by a value of >10% cells with aberration. The total incidence of cells with aberration in the ethyl acetate treated group was only 11%.<sup>16</sup> Hence, based on these results of the single chromosomal aberration assay, it is difficult to interpret that ethyl acetate is mutagenic.<sup>22</sup> Single oral or intraperitoneal administration of ethyl acetate did not damage the bone marrow chromosomes of mice or hamsters as measured by the micronucleus assay.<sup>23,24</sup>

Ethyl acetate is rapidly hydrolyzed to ethyl alcohol in rats, rabbits and human blood. *In vivo* ethyl acetate is rapidly metabolized by rats and humans to ethanol and acetic acid, which can then be disposed of by normal metabolic pathways.<sup>25,26</sup> Ethyl acetate was relatively nontoxic when administered orally, dermally or by inhalation to rats, mice, rabbits and guinea pigs.<sup>27-36</sup> Vapors of ethyl acetate were irritating to the eyes and respiratory tract of humans and various laboratory animal species.<sup>27,37-42</sup> In a 90-day inhalation exposure study in male and female rats exposed to 350, 750 and 1500 ppm, a no-observed-effect-level (NOEL) for male rats was not demonstrated. The lowest-observed-effect-level (LOEL) in female rats was 750 ppm, based on reduction in body weight, body weight gain and food efficiency. The demonstrated NOEL in female rats was 350 ppm.<sup>43</sup> A NOEL of 900 mg/kg was reported in a subchronic gavage study with Sprague-Dawley rats.<sup>44</sup> Subchronic exposure of three guinea pigs to 2000 ppm ethyl acetate, 4 hours/day, 6 days/week, for 65 exposures, led to no measurable changes in body weight gains, or in hematological or urinary values.<sup>45</sup>

No carcinogenic effects of ethyl acetate have been reported in humans. Ethyl acetate did not increase the number of lung tumors in mice when administered by the intraperitoneal route in a 24-week carcinogenicity study.<sup>46</sup>

Neither skin irritation nor sensitization was observed in human subjects tested with a nail polish remover containing 16.5% ethyl acetate<sup>47</sup> or other products containing 97% ethyl acetate.<sup>48</sup> When tested in humans, at concentrations of up to 10% in petrolatum and cosmetic formulations, ethyl acetate was a slight irritant but was not a sensitizer.<sup>30,39,49-51</sup> In a clinical study, ethyl acetate, at 6.5% in a nail color, did not produce signs of phototoxicity or photoallergenicity.<sup>52</sup>

## **TOXICITY DATA ON BURNT MATERIAL**

### ***Combustion studies***

As suggested by the purge and trap studies conducted by PM USA,<sup>53</sup> ethyl acetate applied to tobacco would be expected to extensively distill at 100°C. At the higher temperatures used in the pyrolysis studies, the largest peak was identified as ethyl acetate.<sup>54</sup> The results of this analysis suggest that ethyl acetate would not be pyrolyzed and would be delivered to the smoke mostly intact.

### ***Philip Morris ingredient testing program***

Ethyl acetate was part of a PM USA testing program that was designed to evaluate the potential effects of 333 ingredients added to typical commercial blended test cigarettes on selected biological and chemical endpoints.<sup>55-58</sup> Three pairs of test cigarettes were produced, each containing different groups of ingredients. Ethyl acetate was added to two pairs at target levels of 134 ppm, 172 ppm, 402 ppm and 515 ppm. No significant effects were noted in cytotoxicity, mutagenic studies or in respiratory tract endpoints in 90-day rat inhalation studies. In addition, smoke chemistry studies from cigarettes containing a mixture of flavors including ethyl acetate did not significantly alter the smoke chemistry profile compared to control cigarettes. Based on the results of these studies, the authors concluded that these ingredients (including ethyl acetate) added to tobacco do not add significantly to the overall toxicity of cigarettes.

## **CONCLUSION**

Cigarette smoking causes lung cancer, heart disease, emphysema and other serious diseases in smokers. Smokers are far more likely to develop serious diseases, like lung cancer, than non-smokers. There is no "safe" cigarette. Government health warnings about smoking apply to all cigarettes, regardless of the ingredients added, including those containing only tobacco and paper.

Currently, information is only available for tests utilizing ethyl acetate in a mixture of ingredients applied to cigarette tobacco. Studies are ongoing to address the use of ethyl acetate as a single ingredient. Published studies show there is no meaningful difference in the composition or toxicity of smoke from cigarettes with added ingredients (including ethyl acetate) compared to the smoke from cigarettes without added ingredients.<sup>55-63</sup> It is our scientific judgement, based on the best available data, that ethyl acetate used in our cigarettes does not increase the overall toxicity of cigarette smoke.

## References

1. FEMA (1965) Survey of flavoring ingredient usage levels. No. 2414. *Food Technology* 19(2):155.
2. JECFA (1967) *Toxicological evaluation of some flavoring substances and non-nutritive sweetening agents*. F.A.O. Nutrition Meeting Series. World Health Organization, Joint FAO/WHO Expert Committee on Food Additives, Report No. 44A.
3. CoE (2000) *Chemically-defined Flavouring Substances*. Council of Europe Publishing, F-67075 Strasbourg Cedex, France. p.221.
4. Bartley, J.P. and Schwede, A.M. (1989) Production of volatile compounds in ripening kiwi fruit (*Actinidia chinensis*). *Journal of Agriculture and Food Chemistry* 37:1023-1025.
5. Binder, R.G. and Flath, R.A. (1989) Volatile components of pineapple guava. *Journal of Agriculture and Food Chemistry* 37:734-736.
6. Cadwallader, K.R. and Xu, Y. (1994) Analysis of volatile components in fresh grapefruit juice by purge and trap/gas chromatography. *Journal of Agriculture and Food Chemistry* 42:782-784.
7. Chyau, C.; Chen, S. and Wu, C. (1992) Differences of volatile and nonvolatile constituents between mature and ripe guava (*Psidium guajava* Linn) fruits. *Journal of Agriculture and Food Chemistry* 40:846-849.
8. Moshonos, M.G. and Shaw, P.E. (1994) Quantitative determination of 46 volatile constituents in fresh unpasteurized orange juices using dynamic headspace gas chromatography. *Journal of Agriculture and Food Chemistry* 42(7):1525-1528.
9. Wu, C. and Liou, S. (1992) Volatile components of water boiled duck meat and Cantonese style roasted duck. *Journal of Agriculture and Food Chemistry* 40:338-341.
10. De Vincenzi, M.; Badellino, E.; Di Folco, S.; Dracos, A.; Magliola, M.; Stacchini, A.; Stacchini, P. and Silano, V. (1989) A basis for estimation of consumption: literature values for selected food volatiles. Part III. *Food Addit Contam* 6(2):235-267.
11. TNO-CIVO Food Analysis Institute (1993) Volatile Compounds in Food. Quantitative and Qualitative Data, supplement 4 (series). (H. Maarse and C. A. Visscher, Eds.). Netherlands.
12. Isacoff, H. (1979) *Kirk-Othmer Encycl. Chem. Tech.* 3rd Edition. Wiley Interscience, New York, NY. p.163.
13. FDA (1987) Cosmetic product formulation data: ingredients used in each product category. Computer Printout. Washington, D.C. Cited in CIR Expert Panel, 1989.
14. CIR (Cosmetic Ingredient Review) Expert Panel (1989) Final report on the safety assessment of ethyl acetate and butyl acetate. *Journal of the American College of Toxicology* 8(4):681-705.

15. Ames, B.N.; McCann, J. and Yamasaki, E. (1975) Methods for detecting carcinogens and mutagens with the Salmonella/mammalian-microsome mutagenicity test. *Mutat Res* 31(6):347-364.
16. Ishidate, M., Jr.; Sofuni, T.; Yoshikawa, K.; Hayashi, M.; Nohmi, T.; Sawada, M. and Matsuoka, A. (1984) Primary mutagenicity screening of food additives currently used in Japan. *Food Chem Toxicol* 22(8):623-36.
17. Yamaguchi, T. (1985) Stimulating effects of organic solvents on the mutagenicities of sugar degradation compounds. *Agricultural Biology and Chemistry* 49(12):3363-3368.
18. McCann, J. and Ames, B.N. (1976) Detection of carcinogens as mutagens in the Salmonella/microsome test: assay of 300 chemicals: discussion. *Proc Natl Acad Sci U. S. A.* 73(3):950-4.
19. Shirasu, Y.; Moriya, M.; Kato, K.; Furuhashi, A. and Kada, T. (1976) Mutagenicity screening of pesticides in the microbial system. *Mutat Res* 40(1):19-30.
20. Mayer, V.W. and Goin, C.J. (1988) Investigations of aneuploidy-inducing chemical combinations in *Saccharomyces cerevisiae*. *Mutat Res* 201(2):413-21.
21. Mayer, V.W. and Goin, C.J. (1987) Aneuploidy induced by nocodazole or ethyl acetate is suppressed by dimethyl sulfoxide. *Mutat Res* 187(1):31-5.
22. Burdock, G.A. (2000) Safety Assessment of Ethyl Acetate as a Food Ingredient. Burdock Group, Vero Beach, FL. Unpublished Work.
23. Basler, A. (1986) Aneuploidy-inducing chemicals in yeast evaluated by the micronucleus test. *Mutat Res* 174(1):11-13.
24. Hayashi, M.; Kishi, M.; Sofuni, T. and Ishidate, M., Jr. (1988) Micronucleus tests in mice on 39 food additives and eight miscellaneous chemicals. *Food Chem Toxicol* 26(6):487-500.
25. Sax, N.I. (1984) *Dangers properties of Industrial Materials*. 6th Edition. Van Nostrand Reinhold, New York, NY. p.1315.
26. Tambo, S. (1973) Toxicity hazards of paint thinners with particular emphasis on the metabolism and toxicity of acetate esters. *Nichidai Igaku Zasshi* 32(3):349-360.
27. Smyth, H.F., Jr.; Carpenter, C.P.; Weil, C.S.; Pozzani, U.C.; Striegel, J.A. and Nycum, J.S. (1969) Range-finding toxicity data: List VII. *Am Ind Hyg Assoc J* 30(5):470-6.
28. Patty, F. (1963) *Industrial Hygiene and Toxicology*. 2nd Revised Edition. (F. Patty). Vol. 2. Interscience, New York.
29. Kawasaki, Y.; Suzuki, S.; Kobayashi, K.; Tobe, M. and Ikeda, Y. (1975) [Studies on the inhalation toxicity of A-thinner, its principal constituents and chloroform (author's transl)]. *Eisei Shikenjo Hokoku* (93):72-7.
30. BIBRA (1992) Ethyl acetate. Toxicity Profile. The British Industrial Biological Research Association.

31. Kojima, T.; Takeoka, T.; Yashki, M. and Kobayashi, H. (1977) Toxicological study of toluene poisoning by inhalation: Effect of inhalation of toluene and ethyl acetate mixed vapor on death. *Nippon Hoigaku Zasshi* 31(5):280-290.
32. Fassett, D.W. (1963) In *Industrial Hygiene and Toxicology*. 2nd Edition. (F. Patty, Ed.). Interscience, New York.
33. de Ceaurriz, J.C.; Micillino, J.C.; Bonnet, P. and Guenier, J.P. (1981) Sensory irritation caused by various industrial airborne chemicals. *Toxicol Lett* 9(2):137-143.
34. Kane, L.E.; Dombroske, R. and Alarie, Y. (1980) Evaluation of sensory irritation from some common industrial solvents. *Am Ind Hyg Assoc J* 41(6):451-5.
35. Cosmetic Toiletry and Fragrance Association (1987) Submission of unpublished data by CTFA to the Cosmetic Ingredient Review Director. Acute oral toxicity of a nailpolish remover containing 16.5% ethyl acetate. Cited in CIR Expert Panel, 1989.
36. Munch, J.C. (1972) Aliphatic alcohols and alkyl esters: narcotic and lethal potencies to tadpoles and to rabbits. *IMS Ind Med Surg* 41(4):31-3.
37. Carpenter, C.P. and Smyth, H.F.Jr. (1946) Chemical burns of rabbit cornea. *American Journal of Ophthalmology* 29:1363-1372.
38. Cosmetic Toiletry and Fragrance Association (1980) Submission of unpublished data by CTFA to the Cosmetic Ingredient Review Director. Ocular irritation test. Cited in CIR Expert Panel, 1989.
39. Cosmetic Toiletry and Fragrance Association (1976) Submission of unpublished data by CTFA to the Cosmetic Ingredient Review Director. Cumulative irritation test. Cited in CIR Expert Panel, 1989.
40. Kennah, H.E.I.; Hignet, S.; Laux, P.E.; Dorko, J.D. and Barrow, C.S. (1989) An objective procedure for quantitating eye irritation based upon changes of corneal thickness. *Fundam Appl Toxicol* 12(2):258-68.
41. Kennah, H.E.I.; Albuлесcu, D.; Hignet, S. and Barrow, C.S. (1989) A critical evaluation of predicting ocular irritancy potential from an in vitro cytotoxicity assay. *Fundam Appl Toxicol* 12(2):281-90.
42. Nelson, K.W.; Ege, J.F.Jr.; Woodman, L.E. and Silverman, L. (1943) Sensory response to certain industrial solvent vapors. *Journal of Industrial Hygiene and Toxicology* 25(7):282-285.
43. Christoph, G.R. (1997) Subchronic inhalation neurotoxicity study of ethyl acetate in rats. Haskell Laboratory Report No. 454-96.
44. USEPA (1987) Health and environmental effects profile for ethyl acetate. Report No. EPA/600/X-87/028. NTIS PB89120323.
45. Smyth, H.F. and Smyth, H.F.Jr. (1928) Inhalation experiments with certain lacquer solvents. *Journal of Industrial Hygiene* 10(8):261-271.
46. Stoner, G.D.; Shimkin, M.B.; Kniazeff, A.J.; Weisburger, J.H.; Weisburger, E.K. and Gori, G.B. (1973) Test for carcinogenicity of food additives and chemotherapeutic agents by the pulmonary tumor response in strain A mice. *Cancer Res* 33(12):3069-85.

47. Epinette, W.W. (1987) Prophetic patch and in-use testing of a nail polish remover containing 97% ethyl acetate. Personal communication to the Cosmetic, Toiletry and Fragrance Association. Cited in CIR Expert Panel, 1989.
48. Cosmetic Toiletry and Fragrance Association (1983) The determination of the contact sensitizing potential of a product containing 97% ethyl acetate by means of maximization test. Ivy Research Laboratories, Inc. Cited in CIR Expert Panel, 1989.
49. Hill Top Research Inc. (1984) Submission of unpublished data to the Cosmetic Ingredient Review Director. Report of a human cumulative skin irritation test of a product containing 97% ethyl acetate. Cited in CIR Expert Panel, 1989.
50. Cosmetic Toiletry and Fragrance Association (1980) Submission of unpublished data by CTFA to the Cosmetic Ingredient Review Director. Clinical repeated insult patch test. Cited in CIR Expert Panel, 1989.
51. Cosmetic Toiletry and Fragrance Association (1976) Submission of unpublished data by CTFA to the Cosmetic Ingredient Review Director. Maximization test. Cited in CIR Expert Panel, 1989.
52. Cosmetic Toiletry and Fragrance Association (1980) Submission of unpublished data by CTFA to the Cosmetic Ingredient Review Director. Clinical photopatch test. Cited in CIR Expert Panel, 1989.
53. PM USA (2002) P&T/GC/MS Analysis of ethyl acetate. Request 20020612. Scan TD152HCB. Unpublished Internal Report.
54. PM USA (2002) Pyrolysis GC/MS analysis of ethyl acetate. Request 20020612. Scan P020612A.D. Unpublished Internal Report.
55. Carmines, E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 1: Cigarette design, testing approach, and review of results. *Food and Chemical Toxicology* 40:77-91.
56. Roemer, E.; Tewes, F.J.; Meisgen, T.J.; Veltel, D. and Carmines, E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 3: *In vitro* genotoxicity and cytotoxicity. *Food and Chemical Toxicology* 40:105-111.
57. Rustemeier, K.; Stabbert, R.; Haussmann, H.J.; Roemer, E. and Carmines E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 2: Chemical composition of mainstream smoke. *Food and Chemical Toxicology* 40:93-104.
58. Vanscheeuwijck, P.M.; Teredesai, A.; Terpstra, P.M.; Verbeek, J.; Kuhl, P.; Gerstenberg, B.; Gebel, S. and Carmines E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 4: Subchronic inhalation toxicity. *Food and Chemical Toxicology* 40:113-131.
59. Gaworski, C.L.; Dozier, M.M.; Heck, J.D.; Gerhart, J.M.; Rajendran, N.; David, R.M.; Brennecke, L.H. and Morrissey, R. (1998) Toxicologic evaluation of flavor ingredients added to cigarette tobacco: 13 week inhalation exposures in rats. *Inhal. Toxicol.* 10:357-381.
60. Gaworski, C.L.; Heck, J.D.; Bennett, M.B. and Wenk, M.L. (1999) Toxicologic evaluation of flavor ingredients added to cigarette tobacco: skin painting bioassay of cigarette smoke condensate in SENCAR mice. *Toxicology* 139(1-2):1-17.

61. Doull, J.; Frawley, J.P.; George, W.J.; Loomis, T.A.; Squire, R.A. and Taylor, S.L. (1994) A safety assessment of ingredients added to tobacco in the manufacturing of cigarettes. Covington and Burling, Washington, D.C.
62. Doull, J.; Frawley, J.P.; George, W.J.; Loomis, T.A.; Squire, R.A. and Taylor, S.L. (1998) A safety assessment of ingredients added to tobacco in the manufacturing of cigarettes. Covington and Burling, Washington, D.C.
63. Baker, R.R.; Massey, E.D. and Smith, G. (2004) An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food and Chemical Toxicology* 42S(Supplement 1):S53-S83.